

# NO PAWS ON THE GROUND



We require that all puppies must have at least two DHPP vaccines before they can have “paws on the ground”. Please note that this protocol applies to fosters’ own homes/yards. Public or high traffic/risk areas such as parks, dog parks, animal clinics, etc., should be avoided until they are five months old.

If they do not, they are considered, “NO PAWS ON THE GROUND”. This means that we do not allow these pups to be on the ground outside, including private backyards.

Canine Parvovirus is quite common in San Antonio and can survive in the soil for up to a year or longer all while being resistant to temperature changes. It can travel unknowingly by outside animals (squirrels, raccoons, opossums, etc.) that stepped into an infected area and then traveled into your yard. This hardy virus can also travel well on shoes and clothing that have been in contact with it.

We ask that you read this information thoroughly and help us keep your foster the healthiest it can be!

If think your pup is displaying signs of parvo please contact our Marbach clinic immediately at:

[appointment@sanantoniopetsalive.org](mailto:appointment@sanantoniopetsalive.org).

# What Every Owner Should Know About Parvo in Dogs

By [Anna Burke](#)

Parvo is every new puppy and dog owner's worst nightmare. In a matter of days, a perfectly healthy puppy can go from playful and active to fatally ill. Parvo in dogs is a preventable disease, however, all new puppy owners and breeders need to be aware of the risks of parvo, how to prevent it, and what to do if a puppy catches the parvovirus.

## What Is Parvo?

Parvo is a highly contagious virus. It causes an infectious gastrointestinal (GI) illness in puppies and young dogs, and without treatment, it is potentially deadly.

Part of what makes the virus so dangerous is the ease with which it is spread through the canine population. The virus spreads either by direct contact with an infected dog, or through feces, and an infected dog can begin shedding the virus four-to-five days after exposure — often before the dog starts exhibiting any clinical signs of infection. The dog will continue to shed the virus while he is sick and for up to 10 days after he has recovered. This means that accurate diagnosis and quarantine are essential for the health of your dog and of other dogs, as well.

## What Dogs Are Most at Risk for Parvo?

Young dogs between six weeks and six months old, unvaccinated, or incompletely vaccinated dogs are most at risk for contracting parvo. German Shepherd Dogs, Rottweilers, Doberman Pinschers, English Springer Spaniels, and American Staffordshire Terriers also have an increased risk of contracting the parvovirus, although scientists are not entirely sure why these dog breeds are at a higher risk than others.

Puppies are born with antibodies from their mothers. As these antibodies fade, however, it is up to owners to make sure that the puppies receive a course of parvo vaccinations. The stress of weaning and a secondary parasite or infection, along with parvo, can lead to a more severe case of parvo, which is why it is very important to talk to your vet about the proper care for puppies and pregnant bitches.

## What Causes Parvo in Dogs?

The canine parvovirus causes parvo in dogs, and it can be transmitted in two ways. The first is by direct contact through the nose and mouth with infected poop, which can happen when a dog sniffs or licks a surface or another dog that has been contaminated with feces. Since puppies explore their world through smell and love to mouth things, it is easy to see how a curious puppy could contract the parvovirus.

The second method of transmission is through indirect contact. The virus can survive on clothing, equipment, fur, on human skin, and in the environment. Indirect transmission occurs when a puppy comes into contact with a contaminated person, object, or environment.

The parvovirus is a particularly resilient virus. It can survive indoors at room temperature for at least two months and is resistant to many commonly used cleaners and disinfectants. Outdoors, the parvovirus can survive for months, and even years, if protected from direct sunlight. This is why hospital quarantine of the infected dog and proper cleanup of the environment are especially important.

Shoes that have come into contact with infected feces can also bring the virus into a dog's environment, which is concerning since there is evidence that parvo can live in ground soil for up to one year. If you suspect that you have come into contact with feces at all, you will need to wash the affected area with household bleach, one of the few disinfectants known to kill the virus.

Once a dog has contracted parvo, the virus replicates. This replication takes place in the small intestines, lymphopoietic tissue (lymph nodes, thymus, etc.), and bone marrow. This leads to severe GI problems and in rare cases, myocarditis (inflammation of the heart).

## Symptoms of Parvo in Dogs

Every dog owner and breeder should know the symptoms of parvo in dogs. The most common symptoms are:

- Severe, bloody diarrhea
- Lethargy
- Anorexia
- Fever
- Vomiting
- Weight loss

- Weakness
- Depression
- Dehydration

Any or all of these symptoms merit a call to your veterinarian. Even if parvo is not the cause, the symptoms could be the result of another illness that requires veterinary attention.

## Parvo Prevention

Parvo is a preventable disease, but even vaccinated dogs are not 100% protected from the virus. Vaccines for the parvovirus are recommended for all puppies and are usually given in a series of three shots when the pup is between 6-to-8 weeks old, again at 10-to-12 weeks, and at 14-to-16 weeks. A booster shot is administered one year later and every 3 years after that.

Unvaccinated puppies and incompletely vaccinated puppies should not be exposed to unvaccinated dogs or to environments where unvaccinated dogs could have introduced the parvo virus, like dog parks or boarding facilities. While it might be tempting to take your new puppy with you everywhere you go, her health depends on keeping her safe until she is fully vaccinated against this life-threatening disease.

Understanding parvo in dogs is the first step toward preventing the spread of this dangerous virus. Make sure your puppy gets vaccinated at the appropriate ages and protect your unvaccinated and partially vaccinated puppies by keeping them in a safe environment.